

# AN EXPONENTIAL DIOPHANTINE EQUATION $x^2 + 3^\alpha 113^\beta = y^n$

S.MUTHUVEL AND R.VENKATRAMAN

**ABSTRACT.** The objective of the paper is to determine the complete solutions for the Diophantine equation  $x^2 + 3^\alpha 113^\beta = y^n$  in positive integers  $x$  and  $y$  (where  $x, y \geq 1$ ), non-negative exponents  $\alpha$  and  $\beta$ , and an integer  $n \geq 3$ , subject to the condition  $\gcd(x, y) = 1$ .

## 1. INTRODUCTION

Consider the Diophantine equation of the form

$$(1.1) \quad x^2 + \mathcal{C} = y^n, \quad x, y \geq 1, \quad n \geq 3,$$

where  $\mathcal{C}$  is a fixed positive integer. The initial discovery of positive integer solutions for the given equation dates back almost 17 decades ago [23]. It has been proven that the equation invariably has only a finite number of solutions that are positive integers [21]. Early investigations focused on Eq. (1.1) when  $\mathcal{C} = c_0$  is a constant integer [20, 30, 31]. In [16], Cohn presented a solution to equation (1.1) with the stipulation that  $\gcd(x, y) = 1$ , while taking into account the parameter  $\mathcal{C}$  within the interval  $1 \leq \mathcal{C} \leq 100$ , excluding particular  $\mathcal{C}$  values. The work in [28] addressed additional  $\mathcal{C}$  values within the same range, and the remaining values were addressed in [11]. Over the years, researchers have investigated not only instances where  $\mathcal{C} = \mathfrak{p}^k$  with a specific prime number  $\mathfrak{p}$  [1, 2, 3, 4, 6, 15, 27] but also for the case of a general prime number  $\mathfrak{p}$  [7, 9, 22, 32, 38].

Consider a set of primes  $\mathcal{S} = \{\mathfrak{p}_1, \mathfrak{p}_2, \dots, \mathfrak{p}_k\}$ . Recent investigations focus on Eq. (1.1), specifically when  $\mathcal{C}$  is the product of prime powers  $\mathfrak{p}^k$ , with  $\mathfrak{p} \in \mathcal{S}$  for any non-negative integer  $k$  [5, 10, 13, 14, 17, 18, 19, 24, 25, 26, 29, 33, 34, 35, 36, 37]. Additionally, in [39], the authors explored positive integer solutions to Eq. (1.1) in a more general context, considering  $\mathcal{C} = 2^a \mathfrak{p}^b$  for any odd prime  $\mathfrak{p}$ .

---

2020 *Mathematics Subject Classification.* 11D41, 11D61, 11Y50.

*Key words and phrases.* Diophantine equations, Integer solution, S-integers, Lucas sequence, Primitive divisor.

The primary focus of this paper is the Diophantine equation given by

$$(1.2) \quad x^2 + 3^\alpha 113^\beta = y^n, \quad n \geq 3,$$

where  $\alpha, \beta \geq 0$  and  $x, y \geq 1$  with  $\gcd(x, y) = 1$ .

Now, we proceed to establish the subsequent result.

**Theorem 1.1.** *The equation (1.2) has the following solutions:*

$$(x, y, n, \alpha, \beta) = (2, 7, 3, 1, 1), (1232, 115, 3, 3, 1), (23642486, 82375, 3, 9, 1), \\ (46, 13, 3, 4, 0), (10, 7, 3, 5, 0)$$

excluding the case where  $19 \nmid n$ .

## 2. PRELIMINARIES

Consider algebraic integers  $\eta$  and  $\bar{\eta}$ . A Lucas pair, represented as  $(\eta, \bar{\eta})$ , is defined by the conditions that  $\eta + \bar{\eta}$  and  $\eta\bar{\eta}$  are non-zero coprime rational integers, and  $\frac{\eta}{\bar{\eta}}$  is not a root of unity.

The sequences of Lucas numbers are defined in correspondence with any given Lucas pair  $(\eta, \bar{\eta})$  as follows:

$$\mathcal{L}_n(\eta, \bar{\eta}) = \frac{\eta^n - \bar{\eta}^n}{\eta - \bar{\eta}}, \quad n = 0, 1, 2, \dots$$

The existence of primitive divisors for  $\mathcal{L}_n(\eta, \bar{\eta})$  is a crucial concept in the realm of Lucas sequences.

A prime number  $\mathfrak{p}$  is defined as a primitive divisor of  $\mathcal{L}_n(\eta, \bar{\eta})$  if  $\mathfrak{p} \mid \mathcal{L}_n(\eta, \bar{\eta})$  and  $\mathfrak{p} \nmid (\eta - \bar{\eta})^2 \prod_{i=1}^{n-1} \mathcal{L}_i(\eta, \bar{\eta})$  for  $n > 1$ . Moreover, a primitive divisor  $\mathfrak{q}$  of  $\mathcal{L}_n(\eta, \bar{\eta})$  satisfies  $\mathfrak{q} \equiv \left( \frac{(\eta - \bar{\eta})^2}{\mathfrak{q}} \right) \pmod{n}$ , where  $\left( \frac{*}{\mathfrak{q}} \right)$  denotes the Legendre symbol [?].

For  $n > 4$  and  $n \neq 6$ , all  $n$ -th terms of any Lucas sequence  $\mathcal{L}_n(\eta, \bar{\eta})$  possess primitive divisors, except for certain finite values of parameters  $\eta, \bar{\eta}$ , and  $n$  [8].

## 3. PROOF OF THEOREM 1.1

For the cases of  $n = 3$ ,  $n = 4$ , and  $n \geq 5$ , Equation (1.2) will be investigated independently as follows:

**Proposition 3.1.** *If  $n = 3$ , the Eq. (1.2) has the following solutions:*

$$(x, y, \alpha, \beta) = (2, 7, 1, 1), (1232, 115, 3, 1), (23642486, 82375, 9, 1), \\ (46, 13, 4, 0), (10, 7, 5, 0)$$

*Proof.* When  $n = 3$ , represent  $\alpha = 6a_1 + i$  and  $\beta = 6b_1 + j$ , where  $i, j \in \{0, 1, \dots, 5\}$ . Subsequently, Eq. (1.2) takes the form of an elliptic curve

$$L^2 = M^3 - 3^i 113^j$$

where  $L = \frac{x}{3^{3a_1} 113^{3b_1}}$  and  $M = \frac{y}{3^{2a_1} 113^{2b_1}}$ .

Hence, the problem of finding positive integer solutions for Eq. (1.2) is reduced to finding all  $\{3, 113\}$ -integer points on the relevant 36 elliptic curves for every  $i$  and  $j$ . It's crucial to emphasize that for every finite set of prime numbers  $\mathcal{S}$ , an  $\mathcal{S}$ -integer is characterized as a rational number  $\frac{r}{s}$  where  $r$  and  $s > 0$  are relatively prime integers, and any prime factor of  $s$  is a member of the set  $\mathcal{S}$ .

At this point, we utilized the MAGMA function `SIntegralPoints` to locate all  $\mathcal{S}$ -integral points on the given curves. For  $\mathcal{S} = \{3, 113\}$  [12], the identified points are presented below:

$$(M, L, i, j) = (1, 0, 0, 0), (113, 0, 0, 3), (7, 2, 1, 1), (537, 12444, 2, 1), (3, 0, 3, 0), \\ (15, 18, 3, 1), (51, 360, 3, 1), (115, 1232, 3, 1), (303, 5274, 3, 1), \\ \left(\frac{82375}{9}, \frac{23642486}{27}, 3, 1\right), (353103, 209822526, 3, 1), \\ (339, 0, 3, 3), (13, 46, 4, 0), (7, 10, 5, 0)$$

Considering that  $x$  and  $y$  are positive integers with no common factors, it's important to highlight that only five among these points yield a solution for Eq. (1.2). With this, the proof comes to an end.  $\square$

**Proposition 3.2.** *If  $n = 4$ , the Eq. (1.2) has no positive integer solutions.*

*Proof.* Let  $n = 4$ . Initially, express  $\alpha = 4a_1 + i$  and  $\beta = 4b_1 + j$ , where  $i, j \in \{0, 1, 2, 3\}$ . Consequently, Eq. (1.2) takes the form

$$A^2 = B^4 - 3^i 113^j$$

where  $A = \frac{x}{3^{2a_1} 113^{2b_1}}$  and  $B = \frac{y}{3^{a_1} 113^{b_1}}$ .

Identifying all  $\mathcal{S} = \{3, 113\}$ -integral points on the associated 16 quartic curves corresponds to the search for every integer solution of Eq. (1.2).

By employing the `SIntegralLjunggrenPoints`, we successfully determined all  $\mathcal{S}$ -Integral Points on these curves, resulting in

$$(A, B, i, j) = (\mp 1, 0, 0, 0)$$

Given the condition on the values of  $x$  and  $y$ , it is evident that Eq. (1.2) does not possess any solutions.  $\square$

**Proposition 3.3.** *If  $n \geq 5$ , the Eq. (1.2) does not possess any positive integer solutions.*

*Proof.* Suppose that  $n \geq 5$ . If there exists a solution for Eq. (1.2) with  $n = 2^k$  and  $k \geq 3$ , it can be obtained from solutions with  $n = 4$  since  $y^{2^k} = \left(y^{2^{k-2}}\right)^4$ . Consequently, there are no solutions for (1.2) with  $n = 2^k$  and  $k \geq 3$ . Similarly, (1.2) has no solution for  $n = 3^k$  and  $k \geq 2$ . Hence, without loss of generality,  $n$  is an odd prime.

Let's initiate the analysis of the factorization of Eq. (1.2) in the field  $\mathcal{K} = Q(\sqrt{-d})$  as follows

$$(x + \mathfrak{e}\sqrt{-d})(x - \mathfrak{e}\sqrt{-d}) = y^n$$

where  $\mathfrak{e} = 3^a 113^b$  for some integers  $a, b \geq 0$  and  $d \in \{1, 3, 113, 339\}$ .

Assuming that  $y$  is even leads to a contradiction, as  $x$  must be odd according to (1.2), resulting in  $1 + 3^\alpha \equiv 0 \pmod{8}$ .

As a result,  $y$  is an odd integer, and therefore, the ideals formed by  $x + \mathfrak{e}\sqrt{-d}$  and  $x - \mathfrak{e}\sqrt{-d}$  are relatively prime in the field  $\mathcal{K}$ .

The class number  $h(\mathcal{K})$  takes on one of three values: 1, 6, or 8 for the specific choice of  $d$ .

Thus, we can deduce that the greatest common divisor of  $n$  and  $h(\mathcal{K})$  is 1.

Consider, an algebraic integer  $\xi \in \mathcal{K}$  along with units  $u_1$  and  $u_2$  in the ring of algebraic integers of  $\mathcal{K}$ , we can express this as:

$$x + \mathfrak{e}\sqrt{-d} = \xi^n u_1$$

$$x - \mathfrak{e}\sqrt{-d} = \bar{\xi}^n u_2$$

Considering the orders of the multiplicative group of units in the ring of algebraic integers of  $\mathcal{K}$ , which are 2, 4, or 6 based on the value of  $d$ , and noting that these

orders are relatively prime to  $\mathfrak{n}$ , the presence of units  $u_1$  and  $u_2$  in the equations can be eliminated. The units  $u_1$  and  $u_2$  can be integrated into the factors  $\xi^n$  and  $\bar{\xi}^n$ .

Let us consider the two cases separately, specifically when  $d$  belongs to either  $\{1, 113\}$  or  $\{3, 339\}$  based on distinct integral bases for  $\mathcal{O}_K$  as  $\{1, \sqrt{-d}\}$  and  $\{1, \frac{1+\sqrt{-d}}{2}\}$ , respectively. To begin with, let us assume  $d \in \{1, 113\}$ . Consequently,

$$\begin{aligned} x + \mathfrak{e}\sqrt{-d} &= \xi^n = (s + t\sqrt{-d})^n \\ x - \mathfrak{e}\sqrt{-d} &= \bar{\xi}^n = (s - t\sqrt{-d})^n \end{aligned}$$

and  $y = s^2 + dt^2$  for some rational integers  $s$  and  $t$ . By analyzing these equations, we can derive that

$$\mathfrak{e} = \mathcal{L}_n t,$$

where  $\mathcal{L}_n = \frac{\xi^n - \bar{\xi}^n}{\xi - \bar{\xi}}$ . Notably, the sequence  $\mathcal{L}_n$  is a Lucas sequence.

The Lucas sequences without primitive divisors are explicitly enumerated in [8], and it is confirmed that  $\mathcal{L}_n$  doesn't match any of them. Consequently, we delve into the possibility that a primitive divisor may exist for  $\mathcal{L}_n$ . Assume  $\mathfrak{q}$  is any primitive divisor of  $\mathcal{L}_n$ . In this case,  $\mathfrak{q}$  is either 3 or 113. Considering that any primitive divisor is congruent to  $\pm 1$  modulo  $\mathfrak{n}$ , we rule out the possibility  $\mathfrak{q} = 3$  given that  $\mathfrak{n} \geq 5$ . Thus, we continue with  $\mathfrak{q} = 113$ . According to the definition of a primitive divisor,  $\mathfrak{q} \nmid (\xi - \bar{\xi})^2 = -4dt^2$ , indicating that  $d = 1$ . Furthermore, since  $\left(\frac{-4t^2d}{q}\right) = \left(\frac{-1}{113}\right) = 1$ , we deduce that  $113 \equiv 1 \pmod{\mathfrak{n}}$  which implies that  $\mathfrak{n} = 7$ . Hence, we derive the expression:

$$(3.1) \quad t(7s^6 - 35s^4t^2 + 21s^2t^4 - t^6) = 3^a 113^b$$

where  $\alpha = 2\mathfrak{a}$  and  $\beta = 2\mathfrak{b}$ . Given that  $113 \nmid t$  and  $\gcd(s, t) = 1$ , we conclude that either  $t = \pm 1$  or  $t = \pm 3^a$ . If  $t = \pm 1$ , then Eq. (3.1) transforms into the following equations:

$$7s^6 - 35s^4 + 21s^2 - 1 = \pm 3^a 113^b$$

By letting  $\mathfrak{a} = 2a_1 + i$  and  $\mathfrak{b} = 2b_1 + j$  for  $i, j \in \{0, 1\}$ , we obtain:

$$7s^6 - 35s^4 + 21s^2 - 1 = \delta Y^2, \text{ where } \delta = \pm 3^i 113^j \text{ and } Y = 3^{a_1} 113^{b_1}.$$

By multiplying  $7^2 \delta^3$  both sides of the above equation, we obtain:

$$V^3 - 35\delta V^2 + 147\delta^2 V^2 - 49\delta^3 = (7\delta^2 Y)^2, \text{ where } V = 7\delta s^2.$$

We utilize MAGMA to compute all integral points of these elliptic curves for each value of  $\delta$ , and find the points:

$$\begin{aligned}
(V, 7\delta^2 Y) &= (1, 8), (58, 293) \text{ for } \delta = 1, \\
(V, 7\delta^2 Y) &= (226, 12769) \text{ for } \delta = 113, \\
(V, 7\delta^2 Y) &= (7, 56), (91, 56), (3892, 239519) \text{ for } \delta = 3, \\
(V, 7\delta^2 Y) &= (13195, 672728), (13447, 715064) \text{ for } \delta = 339, \\
(V, 7\delta^2 Y) &= (-21, 56), (-5, 8), (0, 7), (7, 56), (39, 344) \text{ for } \delta = -1, \\
(V, 7\delta^2 Y) &= (-41, 568), (1243, 102152) \text{ for } \delta = -113, \\
(V, 7\delta^2 Y) &= (-2147, 102152), (2230, 331171) \text{ for } \delta = -339.
\end{aligned}$$

It can be verified that none of them leads to a solution of Eq. (1.2).

In the case of  $t = \pm 3^a$ , dividing both sides of Eq. (3.1) by  $t^6$  and introducing  $\mathfrak{b} = 2b_1 + j$  for  $j \in \{0, 1\}$  yields:

$$7 \left( \frac{s^2}{t^2} \right)^3 - 35 \left( \frac{s^2}{t^2} \right)^2 + 21 \left( \frac{s^2}{t^2} \right) - 1 = \pm 113^j \left( \frac{113^{b_1}}{t^3} \right)^2.$$

Setting  $\delta = \pm 113^j$ ,  $Y = \frac{113^{b_1}}{t^3}$ , and multiplying both sides by  $7^2 \delta^3$ , we arrive at:

$$V^3 - 35\delta V^2 + 147\delta^2 V - 49\delta^3 = (7\delta^2 Y)^2, \text{ where } V = \frac{7\delta s^2}{t^2}.$$

Utilizing MAGMA, we determined all  $\{3\}$ -Integral Points on the relevant elliptic curves, resulting in the following set of points:

$$\text{For } \delta = 1, (V, 7\delta^2 Y) = (1, 8), (58, 293)$$

$$\text{For } \delta = 113, (V, 7\delta^2 Y) = (226, 12769)$$

$$\text{For } \delta = -1, (V, 7\delta^2 Y) = (-21, 56), \left( \frac{-77}{9}, \frac{728}{27} \right), (-5, 8), (0, 7), (7, 56), (39, 344)$$

$$\text{For } \delta = -113, (V, 7\delta^2 Y) = (-41, 568), (1243, 102152)$$

Equation (1.2) does not find a solution through any of these cases.

Now, consider the assumption that  $d$  belongs to the set  $\{3, 339\}$ . In such instances, the integral basis for  $\mathcal{O}_K$  is  $\{1, \frac{1+\sqrt{-d}}{2}\}$ , and as a result, we express

$$x + \mathfrak{e}\sqrt{-d} = \xi^n = \left( \frac{s + t\sqrt{-d}}{2} \right)^n$$

$$x - \epsilon\sqrt{-d} = \bar{\xi}^n = \left( \frac{s - t\sqrt{-d}}{2} \right)^n$$

with  $s \equiv t \pmod{2}$ . Then

$$2\epsilon = \mathcal{L}_n t,$$

where  $\mathcal{L}_n = \frac{\xi^n - \bar{\xi}^n}{\xi - \bar{\xi}}$ .

Observing that  $\mathcal{L}_n$  is a Lucas sequence, if there is no primitive divisor for  $\mathcal{L}_n$ , it implies that  $\mathcal{L}_n$  corresponds to one of the Lucas sequences outlined in the table in [8]. However, this observation does not hold true for  $d \in \{3, 339\}$ .

Hence,  $\mathcal{L}_n$  possesses a primitive divisor, denoted as  $\mathfrak{q}$ , where  $\mathfrak{q}$  can either be 3 or 113.

From the properties  $\mathfrak{q} \nmid (\xi - \bar{\xi})^2 = -dt^2$  and  $\mathfrak{q} \equiv \left( \frac{-t^2 d}{\mathfrak{q}} \right) \pmod{n}$ , it follows that  $\mathfrak{q} = 113$ ,  $d = 3$ , and  $n = 19$ . This contradicts the assumption that  $19 \nmid n$ . This completes the proof.  $\square$

## REFERENCES

- [1] F.S. Abu Muriefah, S.A. Arif, The Diophantine equation  $x^2 + 3^m = y^n$ . *Int. J. Math. Math. Sci.* 21(3), 619–620 (1998)
- [2] F.S. Abu Muriefah, S.A. Arif, The Diophantine equation  $x^2 + 5^{2k+1} = y^n$ . *Indian J. Pure Appl. Math.* 30(3), 229–231 (1999)
- [3] F.S. Abu Muriefah, S.A. Arif, On the Diophantine equation  $x^2 + 2^k = y^n$ . *Int. J. Math. Math. Sci.* 20(2), 299–304 (1997)
- [4] F.S. Abu Muriefah, On the Diophantine equation  $x^2 + 5^{2k} = y^n$ . *Demonstratio Math.* 39(2), 285–289 (2006)
- [5] M. Alan, U. Zengin, On the Diophantine equation  $x^2 + 3^a 41^b = y^n$ . *Period. Math. Hung.* 81, 284–291 (2020)
- [6] S.A. Arif, F.S. Abu Muriefah, On the Diophantine equation  $x^2 + 2^k = y^n$  II. *Arab J. Math. Sci.* 7(1), 67–71 (2001)
- [7] S.A. Arif, F.S. Abu Muriefah, On the Diophantine equation  $x^2 + q^{2k+1} = y^n$ . *J. Number Theory* 95, 95–100 (2002)
- [8] Y. Bilu, G. Hanrot, P.M. Voutier, Existence of primitive divisors of Lucas and Lehmer numbers (with Appendix by Mignotte). *J. Reine Angew. Math.* 539, 75–122 (2001)
- [9] A. Bérczes, I. Pink, On the Diophantine equation  $x^2 + p^{2k} = y^n$ . *Arch. Math.* 91, 505–517 (2008)
- [10] A. Bérczes, I. Pink, On generalized Lebesgue–Ramanujan–Nagell equation. *An. St. Univ. Ovid. Cons.* 22(1), 51–57 (2014)
- [11] Y. Bugeaud, M. Mignotte, S. Siksek, Classical and modular approaches to exponential Diophantine equations II. The Lebesgue–Nagell equation. *Compos. Math.* 142, 31–62 (2006)
- [12] W. Bosma, J. Cannon, C. Playoust, The Magma algebra system. I. The user language. *J. Symbolic Comput.* 24(3), 235–265 (1997)

- [13] I.N. Cangúl, M. Demirci, F. Luca, Á. Pintér, G. Soydan, On the Diophantine equation  $x^2 + 2^a 11^b = y^n$ . *Fibonacci Q.* 48(1), 39–46 (2010)
- [14] I.N. Cangúl, M. Demirci, M. Inam, F. Luca, G. Soydan, On the Diophantine equation  $x^2 + 2^a 3^b 11^c = y^n$ . *Math. Slovaca* 63(3), 647–659 (2013)
- [15] J.H.E. Cohn, The Diophantine equation  $x^2 + 2^k = y^n$ . *Arch. Math. Basel* 59(4), 341–344 (1992)
- [16] J.H.E. Cohn, The Diophantine equation  $x^2 + C = y^n$ . *Acta Arith.* 65(4), 367–381 (1993)
- [17] M. Demirci, On the Diophantine equation  $x^2 + 5^a p^b = y^n$ . *Filomat* 31(16), 5263–5269 (2017)
- [18] H. Godinho, D. Marques, A. Togbé, On the Diophantine equation  $x^2 + 2^\alpha 5^\beta 17^\gamma = y^n$ . *Commun. Math.* 20(2), 81–88 (2012)
- [19] H. Godinho, D. Marques, A. Togbé, On the Diophantine equation  $x^2 + C = y^n$  for  $C = 2^a 3^b 17^c$  and  $C = 2^a 13^b 17^c$ . *Math. Slovaca* 66(4), 565–574 (2016)
- [20] C. Ko, On the Diophantine equation  $x^2 = y^n + 1$ ,  $xy \neq 0$ . *Sci. Sin.* 14, 457–460 (1965)
- [21] F. Landau, A. Ostrowski, On the Diophantine equation  $ay^2 + by + c = dx^n$ . *Proc. Lond. Math. Soc.* 19(2), 276–280 (1920)
- [22] M. Le, An exponential Diophantine equation. *Bull. Austral. Math. Soc.* 64, 99–105 (2001)
- [23] L.A. Lebesgue, Sur l'impossibilité, en nombres entiers, de l'équation  $x^m = y^2 + 1$ . *Nouv. Ann. Math.* 9(1), 178–181 (1850)
- [24] F. Luca, On the equation  $x^2 + 2^a 3^b = y^n$ . *Int. J. Math. Math. Sci.* 29(3), 239–244 (2002)
- [25] F. Luca, A. Togbé, On the equation  $x^2 + 2^a 5^b = y^n$ . *Int. J. Number Theory* 4(6), 973–979 (2008)
- [26] F. Luca, A. Togbé, On the equation  $x^2 + 2^\alpha 13^\beta = y^n$ . *Colloq. Math.* 116(1), 139–146 (2009)
- [27] F. Luca, A. Togbé, On the Diophantine equation  $x^2 + 7^{2k} = y^n$ . *Fibonacci Quart.* 45(4), 322–326 (2007)
- [28] M. Mignotte, B.M.M. DeWeger, On the Diophantine equations  $x^2 + 74 = y^5$  and  $x^2 + 86 = y^5$ . *Glasgow Math. J.* 38(1), 77–85 (1996)
- [29] S. Muthuvel, R. Venkatraman, An Exponential Diophantine Equation  $x^2 + 3^a 97^b = y^n$ . *Int. J. Math. Comput. Sci.* 19(4), 1125–1128 (2024)
- [30] T. Nagell, Sur l'impossibilité en nombres entiers de quelques équations a deux indéterminés. *Norsk. Mat. Forenings Skifter Ser. I* 13(1), 65–82 (1923)
- [31] T. Nagell, Contributions to the theory of a category of Diophantine equations of the second degree with two unknowns. *Nova Acta Reg. Soc. Upsal. IV Ser.* 2(15), 1–38 (1955)
- [32] X. Pan, The exponential Lebesgue-Nagell equation  $x^2 + p^{2m} = y^n$ . *Period. Math. Hung.* 67(2), 231–242 (2013)
- [33] I. Pink, On the Diophantine equation  $x^2 + 2^a 3^b 5^c 7^d = y^n$ . *Publ. Math. Debrecen* 70(1–2), 149–166 (2007)
- [34] I. Pink, Z. Rabai, On the Diophantine equation  $x^2 + 5^k 17^l = y^n$ . *Commun. Math.* 19, 1–9 (2011)
- [35] S. G. Rayaguru, On the Diophantine equation  $x^2 + C = y^n$ . *Indian J. Pure Appl. Math.*, 1–9 2022
- [36] G. Soydan, M. Ulas, H. Zhu, On the Diophantine equation  $x^2 + 2^a 19^b = y^n$ . *Indian J. Pure Appl. Math.* 43(3), 251–261 (2012)
- [37] G. Soydan, N. Tzanakis, Complete solution of the Diophantine equation  $x^2 + 5^a 11^b = y^n$ . *Bull. Hellenic Math Soc.* 60, 125–152 (2016)
- [38] H. Zhu, A note on the Diophantine equation  $x^2 + q^m = y^3$ . *Acta Arith.* 146(2), 195–202 (2011)



- [39] H. Zhu, M. Le, G. Soydan, A. Togbé, On the exponential Diophantine equation  $x^2 + 2^a p^b = y^n$ . *Period. Math. Hung.* 70, 233–247 (2015)

DEPARTMENT OF MATHEMATICS, FACULTY OF ENGINEERING AND TECHNOLOGY, SRM INSTITUTE OF SCIENCE AND TECHNOLOGY, VADAPALANI CAMPUS, No.1 JAWAHARLAL NEHRU SALAI, VADAPALANI, CHENNAI-600026, TAMIL NADU, INDIA.

*Email address:* muthushan15@gmail.com, ms3081@srmist.edu.in

DEPARTMENT OF MATHEMATICS, FACULTY OF ENGINEERING AND TECHNOLOGY, SRM INSTITUTE OF SCIENCE AND TECHNOLOGY, VADAPALANI CAMPUS, No.1 JAWAHARLAL NEHRU SALAI, VADAPALANI, CHENNAI-600026, TAMIL NADU, INDIA.

*Email address:* venkatrr1@srmist.edu.in